

Fungicide resistance management in corn, soybean, and wheat in Wisconsin

Authors: Paul Esker, Assistant Professor and Extension Field Crops Plant Pathologist, UW-Madison and UW-Extension and Richard Proost, Regional Agronomist, Nutrient and Pest Management Program, UW-Madison

FRAC CODE	GROUP NAME	CHEMICAL FAMILY	ACTIVE INGREDIENT	PATHOGEN RESISTANCE?	MANUFACTURER	PRODUCT EXAMPLE	REGISTERED FOR CROP*
1	MBC's methyl benzimidazole carbamates	thiophanates	thiophanate-methyl	Yes	Cerexagri	<i>Topsin® M 4.5FL</i> <i>Topsin® M 70WP</i> <i>Topsin® M WDG</i>	Soybean, Wheat Soybean, Wheat Soybean, Wheat
3	DMI's demethylation inhibitors	triazole	metconazole	Yes	BASF	<i>Caramba™</i>	Soybean, Wheat
			fenbuconazole	Yes	Dow AgroSciences	<i>Indar®</i>	Wheat
			tetraconazole	Yes	Valent	<i>Domark® 230 ME</i>	Soybean
			tebuconazole	Yes	Bayer CropScience	<i>Folicur®</i>	Wheat
			propiconazole	Yes	Syngenta	<i>Tilt®</i>	Corn, Soybean, Wheat
				Yes	Dow AgroSciences	<i>PropiMax® EC</i>	Soybean, Wheat
			prothioconazole	Yes	Bayer CropScience	<i>Proline® 480SC</i>	Wheat
7	Carboxamides	pyridine carboxamides	boscalid	Yes	BASF	<i>Endura®</i>	Soybean
11	Qol's Quinone outside inhibitors	methoxycarbamate methoxyacrylate	pyraclostrobin azoxystrobin	Yes Yes	BASF Syngenta	<i>Headline®</i> <i>Quadris®</i>	Corn, Soybean, Wheat Corn, Soybean, Wheat
M	Multi-Site Contact Activity	chloronitriles	chlorothalonil	No	Syngenta Sipcam Agro USA	<i>Bravo WeatherStik®</i> <i>Echo® 720</i> <i>Echo® 90DF</i> <i>Echo® Zn</i>	Soybean Soybean Soybean Soybean
		dithiocarbamate	mancozeb	No	Dow AgroSciences DuPont Cerexagri	<i>Dithane® DF Rainshield™</i> <i>Dithane® F-45 Rainshield™</i> <i>Dithane® M45</i> <i>Manzate® ProStick™</i> <i>Manzate® Flowable™</i> <i>Penncozeb™ 75DF</i> <i>Penncozeb™ 80WP</i> <i>Penncozeb™ 4FL</i>	Corn, Wheat Corn, Wheat Corn, Wheat Corn, Wheat Corn, Wheat Corn, Wheat Corn, Wheat Corn, Wheat Corn, Wheat
		inorganic	copper sulfate	No		<i>Cuprofix® MZ Disperss®</i> <i>Cuprofix Ultra 40</i>	Soybean, Wheat Soybean, Wheat
PREMIX FUNGICIDES							
3	DMI	triazole	propiconazole	Yes	Syngenta	<i>Quilt®</i>	Corn, Soybean, Wheat
11	Qol	methoxyacrylate	azoxystrobin	Yes			
3	DMI	triazole	propiconazole	Yes	Bayer CropScience	<i>Stratego®</i>	Corn, Soybean, Wheat
11	Qol	oximinoacetate	trifloxystrobin	Yes			

* MAY BE REGISTERED ON CROPS OTHER THAN CORN, SOYBEAN, OR WHEAT. SEE LABEL FOR INFORMATION.

FRAC and the FRAC code

The Fungicide Resistance Action Committee (FRAC) is an organization made up of representatives of the agro-chemical industry. Their mission: "... is to provide fungicide resistance management guidelines to prolong the effectiveness of 'at risk' fungicides and to limit crop losses should resistance occur." FRAC developed a code, known as the FRAC Code, that can be used to classify fungicides into groups based on their mode of action. Fungicides currently registered in Wisconsin for use on corn, soybean, and wheat fall into five FRAC Codes: 1, 3, 7, 11, and M.

FRAC Code 1: The methyl benzimidazole carbamate (MBC) fungicide group contains the benzimidazole and thiophanate fungicide families. These fungicides are effective against a broad range of fungi that cause leaf spots, root and crown rots, stem rots, and powdery mildews, but not rusts. MBC fungicides inhibit tubulin production, interfering with normal cell division in sensitive fungi. These fungicides have preventative and early-infection activity. While they have systemic properties, they cannot move down the plant, making canopy penetration and complete plant coverage essential for control.

The MBC fungicide risk of resistance is high. The modification of a single amino acid in a fungus can result in resistance. Resistance to these fungicides was first reported in 1970. Many important fungal plant pathogens have become resistant to these fungicides.

FRAC Code 3: The demethylation inhibitors (DMI) fungicide group contains the triazole fungicides. DMI fungicides are highly effective against powdery mildews, rusts, and many leaf spotting fungi. These fungicides work by inhibiting a specific enzyme that plays a role in sterol production in fungi. Sterols are necessary for the development of functional cell walls. Application of DMIs results in abnormal fungal growth and death. However, triazoles have no effect on spore germination because spores contain enough sterol for the formation of germ tubes. Thus, DMI fungicide must be applied preventively or at early-infection to be effective. DMI fungicides are locally systemic, meaning that they are taken up by the leaves and move within the leaf but not necessarily from one leaf to another. Generally, these fungicides have approximately 14 days of residual activity.

DMI fungicides have a very specific site of action so risk of

resistance development is a concern. Resistance management practices include avoiding repeated applications of DMI fungicides alone in the same season against high-risk pathogens such as powdery mildew.

FRAC Code 7: Carboxamide fungicides include boscalid, carboxin, and flutolanil. Boscalid is primarily a foliar fungicide used against the *Botrytis*, *Sclerotinia*, and *Alternaria* pathogens. They work by inhibiting the respiration of target fungi, specifically the complex II fungal respiration. Carboxamide fungicides have locally systemic to systemic properties and movement is translaminar. Resistance has been documented for these fungicides.

FRAC Code 11: The Quinone outside inhibitors (Qol) fungicide group contains three fungicide families, strobilurins, imidazoles, and oxazoles. Qol fungicides are very effective against a broad spectrum of fungi. These fungicides work by inhibiting mitochondrial respiration, effectively stopping energy production of the fungus, and result in death. These fungicides are effective on spore germination and early growth. However, they are not effective against fungi that are growing inside the leaf tissue so they must be applied preventively or at early-infection to be effective. Similar to DMI fungicides, they are locally systemic. These fungicides have approximately 7–21 days of residual activity.

Qol fungicides have a very specific site of action so the risk of resistance development is a high. Currently there are more than 20 plant pathogens with some level of resistance to Qol fungicides.

FRAC Code M: Multi-site activity fungicides include inorganic compounds, dithiocarbamates, and chloronitriles. Multi-site activity fungicides have a broad spectrum of disease control activity. They are considered protective or preventative fungicides as they are applied to the leaf and stem surfaces prior to pathogen appearance. They do not affect fungi once they have infected the plant. Multi-site activity fungicides affect multiple biochemical sites in fungi, killing fungi by overwhelming them with toxins. These fungicides are sensitive to rainfall and sunlight since they are not absorbed into the plant, generally remaining active for 7–14 days.

Multi-site activity fungicides have a low risk of resistance development. Because of this, multi-site activity fungicides are an important part of fungicide resistance management. When multi-site fungicides are combined with either a code 3 or 11 fungicide (if allowed by the

Guidelines for fungicide resistance management

- Plant disease resistant hybrids/varieties whenever possible.
- Scout fields on a regular basis noting incidence and severity of diseases. Use this information to develop a field history for future disease management decisions.
- Tank mix high-risk fungicides with fungicides that have different modes of action, are active against the targeted disease(s), and have similar lengths of residual activity.
- Do not use reduced rates of fungicides.
- Alternate or tank mix fungicides with different modes of action when multiple applications are required.
- Apply fungicides preventively or early in the disease cycle and when a disease threat is warranted.
- Avoid curative fungicide applications, especially with high-risk fungicides.

Always read and follow the pesticide label

- for maximum number of sprays per season.
- for recommended application rates.
- for application timing for both target disease and plant growth stage.

fungicide label), they may extend the number of years those higher risk fungicides can be used by reducing the number of applications of those high-risk fungicides.

Text adapted from:

Mueller, Daren S., and Carl A. Bradley. 2008. Field Crop Fungicides for the North Central United States. Ames, IA and Urbana-Champaign, IL: Iowa State University and University of Illinois North Central Integrated Pest Management Center.

The Fungicide Resistance Action Committee provides a list of fungicides sorted by modes of action: www.frac.info/frac/index.htm.

Copyright © 2009 by the Board of Regents of the University of Wisconsin System doing business as the division of Cooperative Extension of the University of Wisconsin-Extension. All rights reserved. Send copyright inquiries to: Cooperative Extension Publishing, 432 N. Lake St., Rm. 227, Madison, WI 53706.

Authors: **Paul Esker**, Assistant Professor and Extension Field Crops Plant Pathologist, UW-Madison and UW-Extension and **Richard Proost**, Regional Agronomist, Nutrient and Pest Management Program, UW-Madison. Cooperative Extension publications are subject to peer review.

University of Wisconsin-Extension, Cooperative Extension, in cooperation with the U.S. Department of Agriculture and Wisconsin counties, publishes this information to further the purpose of the May 8 and June 30, 1914 Acts of Congress. An EEO/AA employer, the University of Wisconsin-Extension, Cooperative Extension provides equal opportunities in employment and programming, including Title IX and Americans with Disabilities (ADA) requirements. If you need this information in an alternative format, contact Cooperative Extension Publishing or Equal Opportunity and Diversity Programs, University of Wisconsin-Extension, 501 Extension Building, 432 N. Lake Street, Madison, WI 53706, diversity@uwex.edu, phone: (608) 262-0277, fax: (608) 262-8404, TTY: 711 Wisconsin Relay.

This publication is available from your Wisconsin county Extension office (www.uwex.edu/ces/cty) or from Cooperative Extension Publishing. To order, call toll-free: 1-877-947-7827 (WIS-PUBS) or visit our web site: learningstore.uwex.edu.

A3878 Fungicide resistance management in corn, soybean, and wheat in Wisconsin **I-04-2009**